Final Scientific Report for

Title: Synthesis of Atropine Labeled with C14 in the Tropine Ring and Rifery

Cold Acclimation on Atropine Metabolism.

## Bibliography:

- Schmidt, G.S., Eling, T.E. and Drack, J.C.: Synthesis of Tropine-Labeled Alexander of Tropine and for its Esterification with Tropic Acid. J. Pharmaceut. Sci. 56: 215 - 221, 1967.
- Kalser, S.C. and Kunig, R.: Effect of Varying Periods of Cold Exposure on the Action and Metabolism of Hexobarbital. Submitted to J. Pharmacol. exp. Ther. in May, 1967.
  - Kalser, S.C., Evans, E., Forbes, E., Kelly, M., Kelvington, E., Kunig, R. and Randolph, M.: Decreased Atropine Toxicity in Rats Chronically Expessed to Cold To be submitted to Toxicology and Applied Pharmacology, August, 1967.
  - Schmidt, G.S. and Eling, T.S.: Progress report on Synthesis of Ring-labeled Atropines. Several papers to be published when syntheses completed.
  - Kalser, S. C. and McLain, P.L.: Tropine-labeled Atropine Metabolism in Man: N-C14H3-Atropine. Progress report, paper to be published when study completed.

Summary of Results of Research accomplished under Grant AFOSR 87:

Synthesis of a series of  $c^{14}$ -tropine-labeled atropines, which have previously been unavailable, has been accomplished. A few of these compounds are still being synthesized, but when the series is completed, it will represent the most thorough analysis of the tropine ring to be undertaken. Every carbon in the ring will have been tagged separately or in various combinations. In addition, the N-Clung-atropine has also been prepared and made available.

We have shown that environmental conditioning to cold can almost double the LD50 for atropine compared with the LD50 of control rats; however, an acute exposure to cold decreases the LD50. These changes in toxicities parallel the rate of absorption of atropine from the peritoneal cavity; the rate of absorption is slowest in the coldacclimated and fastest in the rat exposed to cold acutely. Excretion and metabolism of atropine is not increased in the cold-acclimated rat. This suggests that the rate of absorption of atropine probably modulates the toxicity pattern of the drug.

Using an N-C14H3-atropine, a study in normal man showed that 91% of the i.m. dose was excreted; 88% in the urine in 48 hours and 3 % in the expired air in 3 hours. This suggests that demethylation of atropine occurs as one of the metabolic pathways for

성


## Final Scientific Report for AFOSK 87-66 pg 2

degralation in man. Most of the urinary products appear to be the unchanged drug and tropine, in about equal amounts. However, this phase of the study is still in its preliminary stage. The drug disappears exceedingly rapidly from the blood.

The urinary excretion pattern suggests at least 3 rates of clearance. Since one of these rates may relate to an enterohepatic phase of recirculation of the drug, biliary excretion will be studied in man. At least a few of the ring-labeled atropines will also be studied in man.

To the state of th

The objective of this investigation was to synthesize atropine labeled with  $c^{14}$  in the propine molety. The proposed method for selective Lubeling of the tropine ring of atropine is outlined in the schemes below.



2,3,4.3<sup>14</sup> Citric Acid

2,3-0<sup>14</sup> Citric Acid

3-0<sup>14</sup> Citric Acid

Acetone dicarboxylic acid

Succindialdehyde Methylamine HCl.

Tropinone

Tr. ne

Atropine

Me IIH<sub>2</sub> — HCl — C<sup>14</sup>

Succindialdehyde Acetone dicarboxylic acid

Tropinone

Tropine

Atropine

.

<del>(44, 116, 16, 16)</del>

During the first year, the investigation was directed to scaling down from 5 gm. to 300-400 mg level the reaction sequence (see below) from , succinfo acid to succindialdehyde.

Succinic acid	N-N - Dimethyl	•
♦ .	Succinanilide	Succindial dehyde
		Tropinone

From 354 mg of succinic aicd, tropinone was obtained in yields ranging from 65-85 mg. The percentage yield of tropinone based on succinic aicd averaged approximately 20%.

Detailed studies for the formation of tropinone from precursor, conversion of tropinone to tropine, and formation of atropine via the esterification of tropine with tropic acid was also conducted. This work, published in the Journal of Pharmaceutical Science; 1967, established the reproducibility of these steps which is required for C<sup>14</sup> labeling of the tropine moiety of atropine.

The primary objective of the second year was the labeling of atropine. During this year, problems arose in the reduction of tropinone to atropine which delay labeling. Our original method for reduction was discarted, and a new procedure for reduction developed.

Since the cheapest C<sup>14</sup> starting material was methylamine -C<sup>14</sup> HCl., this labeled intermediate was first used in order to check the entire procedure for possible new problems. After five unsuccessful labeling attempts, H-He<sup>14</sup> atropine was pooled, the C<sup>14</sup> tropinone isolated and converted to

N-Me<sup>14</sup> tropine. This procedure resulted in the recovery of 35 mg of N-Me<sup>14</sup> tropine which could be used in limited metabolic studies and for standards on chromatograms.

Two attempts at labeling atropine are in progress at grant termination. As of this date we have 50 mg of  $3-0^{14}$  tropine and are preparing for esterification with tropic acid to form  $3-0^{14}$  atropine. Also  $2,3,4-0^{14}$  tropinone has been synthesized but the yield has not been determined. This labeled intermediate is in the process of purification before subsequent conversion to  $2,3,4-0^{14}$  atropine.

We feel that the synthesis has reached the point where it is just a matter of "grinding our" the C<sup>14</sup> labeled compounds. The method from citric acid appears to be working and the method from succinic acid, using cold starting materials, has worked well. The following labeled atropines are expected to be available in a few months:

Labeled Intermediate	Position of Label in Atropine					
3-0 <sup>14</sup> citric acid	3-ć <sup>14</sup>					
2,4-C <sup>14</sup> citric acid	2,4-0 <sup>14</sup>					
2,3,4-0 <sup>14</sup> citric acid	2,3,4-C <sup>14</sup>					
2,3-C <sup>14</sup> succinc acid	6,7-c <sup>14</sup>					
1,4-C14 succinic aicd	1,5-0 <sup>14</sup>					

Two manuscripts are in preparation, the first concerning the prototype migro method for tropine labeling of atropine from arabinose and the second concerning the synthesis of N-Me  $C^{1/4}$  atropine.

Once N-Mc-C<sup>14</sup>-atropine was available, metabolic studies with the labeled drug, in mice, was initiated. At a dose of 100 mg/kilo, approximately 10% of the injected trug was metabolized to C<sup>14</sup>O<sub>2</sub> which was not found in previous metabolic studies. This indicates that one possible metabolite is N-demethyl atropine or one of its derivatives. Chromatographic analysis, of the urine, revealed six major peaks. Of the injected dose, only 50 percent was recovered in expired air, urine, and feces. Since the remaining 50 percent of injected C<sup>14</sup> is mostly in the animals, method are being sought to determine the C<sup>14</sup> remaining in the intact animals. Chromatographic analysis of the urine of mice receiving N-Me<sup>14</sup> tropine in a dose of 180 mg/kilo revealed one major peak at Rf-.55 and two minor peaks.

In an effort to identify the tropine modified metal-clites of atropine, metabolic studies will be conducted with each labeled atropine as they become available:

The state of the s	water to a man an Marchan		The second secon	-	
<b>-#</b>			# · · ·		
Unclassified					
Security Classification					
· , DOCUMENT CO	ONTROL DATA - R&	D		$\neg$	
(Security classification of title, body of abatiscs and title	zing ennotation must be on			_	
- ORIGINATING ACTIVITY (Corporate author)		24. REPORT SECURITY C LASSIFICATION			
Univ. of Pittsburgh, School of Madicin	ne, Pittsburgh,	Unclassified			
Pa.		25 GROUP			
3. REPORT TITLE					
			3 788 6. 6.33	ŀ	
Synthesis of Atropine labeled with Cl4	in the Tropine	ring;	and Filect of Cold	- [	
acclimation on Atropine metabolism.					
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)				-	
Scientific FINAL				- 1	
S. AUTHOR(S) (Last name, liret name, initial)	···			ᅱ	
Schmidt, G.C.				1	
Eling, T.E.				- [	
				I	
6. REPORT DATE	74. TOTAL NO. OF P	AGES	75. NO. OF REFS	可	
July, 1967	<b>8</b> 7		35		
BE. CONTRACT OR GRANT NO. AFOSR 87-166	94. ORIGINATOR'S RI	EPORT NUM	BER(S)	$\neg$	
A PROJECT NO. 9777-01				1	
			~~~		
· 61445014	AL OTHER REPORT	NO(S) (Any	other numbers that may be essione:	1	
1. 681312	AFOSR	67°	2030		
10. AVAILABILITY/LIMITATION NOTICES					
				- [	
Distrib	ution of this				
documen	t is unlimited			- 1	
11. SUPPLEMENTARY NOTES	12. SPONSORING MILI	TARY ACT	VITY	ᅱ	
· \ .			Scientific Research	ı	
TECH, OTHER	1400 Wilson I	oulevar	d (SRLA)	ł	
	Arlington, Vi				
13- ABSTRACT		-0		7	
During the first year, the investigation	ion was directed	to sca	ling down from 5 gm.	to	
300-400 mg level the synthesis of trop	pinone from succ	inic ac	id. Detailed studies	Kol	
the formation of tropinone from precu					
formation of atropine via the esterif					
ducted this work is published in the	Journal of Phar	maceuti	cal Science 56:215,19	57.	
The primary objective of the second y					
problems which arose in the reduction	of tropinone; it	he orig	inal reduction method	1	
was discarded and a new procedure deve a small yield of N-Mell tropine was of prepared from the N-Mell-tropine; this	olopedAfter s	everal	abortive attempts.		
a small yield of N-Meta tropine was of	otained. A batch	of N-M	ed_atropine has been	i	
prepared from the N-Mety-tropine; this	s material has b	een use	d for some metabolic	1	
studies in mice and part has been dist	tributed to S.C.	Kalser :	for metabolic studies		
in humans. Two attempts at making rin	ng-labeled atrop	ino are	in progress. The 3-C		
and the 2,3,4=2-7-atroping have been r	prepared, but th	e yield	s and specific activit	id	
were lower than anticipated. The remains	and the second of the second o				
made in the next few months.	mining o ring	TSDSTEG	atropines will be	- 1	

DD .5084. 1473

Unclassified

Security Classification

## Unclassified

de

Ô

δ

0

· Security Classification

Crawter.

14,0	KEY WORDS	LINK A		LINK.8		LINKC	
سنيا			WT	ROLE	" , WT	ROLE	WT
0 11,2	Cli-ring labeled atropine, synthesis of N-Melt-atropine, synthesis of	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			•		-
	•	ľ l		ļ			-
		], [				.	
ľ	•				<del>.</del>		
,.					•		
, (				,			
	* * * * * * * * * * * * * * * * * * * *	1.	:	,			
-	· .		-		-		
	Q y			I			ļ
ક	. •	.	i	1			
	8	1		]		٠	
	co co			}		ı	
~	. •	ľ	,	l		, ,	;

## INSTRUCTIONS

- 1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.
- 2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations:
- 2b. GRGUP: Automatic dewngrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
- 3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in pare these immediately following the title.
- 4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period in covered.
- 5. AUTHOR(5): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of mervice. The name of the principal author is an absolute minimum requirement.
- G. REPORT DATE: Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.
- 7s. TOTAL NUMBER OF PAGES: The total page count should follow normal pegination procedures, i.e., enter the number of pages containing information.
- 7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.
- 8a. CONTRACT OR GRANT NUMBER. If appropriate, enter the applicable number of the contract or grant under which the report was written.
- 85, &; & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.
- 9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (outher by the originator or by the aponsor), also enter this number(s).
- 10. AVAIDABILITY/LIMITATION NOTICES: Enter any limitations on tuchor dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1). "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign unnouncement and disser. atlon of this report by DDC is not authorized."
- (3) "Ü. S. Government agencies may chisin copies of this report directly from DPC. Other qualified DDC users shall request through
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through

If the report has been furnished to the Office of Technical Servicus, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

- 11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.
- 12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or luboratory sponsoring (paying for) the resourch and development. Include address.
- 13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation shoet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each puragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS):(S). (C). or (U).

There is no limitation on the longth of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

Unclassified

Security Classification